



## EZ-Tn5™ Transposase

Cat. No. TNP92110

Available exclusively thru Lucigen.  
[lucigen.com/epibio](http://lucigen.com/epibio)

**epicentre**<sup>®</sup>  
an Illumina® company

## 1. Introduction

EZ-Tn5™ Transposase\* is a hyperactive, mutated form of Tn5 transposase and a highly efficient enzyme for insertion of an EZ-Tn5 Transposon into any target DNA, *in vitro*.<sup>1</sup> In addition to EZ-Tn5 Transposase, efficient transposition requires that each EZ-Tn5 Transposon have a specific 19-bp transposase recognition sequence (Mosaic End or ME sequence) at each of its ends. Mutations engineered into both the 19-bp MEs and the EZ-Tn5 Transposase results in an *in vitro* transposition frequency that is 1000-fold greater than wild type.

EZ-Tn5 Transposase catalyzes a multi-step “cut and paste” transposition reaction. Initially, the enzyme binds the 19-bp ME of the transposon to form a Transposome™ (synaptic complex). The transposome then randomly attacks and cleaves the phosphodiester backbone of the target DNA. Finally, the EZ-Tn5 Transposase catalyzes the covalent linkage of the 3'-OH ends of the transposon to the exposed 5'-phosphorylated ends of the target DNA. Transposition creates a 9-bp sequence duplication immediately flanking the transposon insertion site.

### EZ-Tn5 Transposase can be used to:

1. Insert a marker, T7 promoter, or R6K $\gamma$  origin of replication flanked by the 19-bp MEs of Lucigen's EZ-Tn5 standard Transposons into any target DNA.
2. Insert any custom DNA sequence flanked by the 19-bp MEs of an EZ-Tn5 Transposon into any target DNA.
3. Prepare Transposomes, in the absence of Mg<sup>2+</sup>, for electroporation into living bacteria and subsequent random insertion of the transposon into the bacterial chromosome.

## 2. EZ-Tn5™ Transposase Components

Component Name	Volume
EZ-Tn5 Transposase	@ 1 U/ $\mu$ l 10 U
EZ-Tn5 10X Reaction Buffer	100 $\mu$ l
EZ-Tn5 10X Stop Solution	100 $\mu$ l
Sterile Water	1 ml

### 3. Product Specifications

**Storage:** Store only at –20°C in a freezer without a defrost cycle.

**Storage Buffer:** EZ-Tn5 Transposase is supplied in a 50% glycerol solution containing 50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1 mM EDTA, 0.1% Triton® X-100, and 1 mM dithiothreitol.

**Unit Definition:** One unit of EZ-Tn5 Transposase catalyzes the release of the donor backbone fragment from 1 µg of transposed DNA in 1 hour at 37°C, as determined by agarose gel electrophoresis.

**Enzyme Structure:** Single polypeptide of 55 Kd.

**Contaminating Activity Assays:** All components of the EZ-Tn5 Transposase are free of detectable DNase and RNase activities as judged by agarose gel electrophoresis following over-digestion assays, with the exception of the inherent endonucleolytic function of the EZ-Tn5 Transposase.

**EZ-Tn5 10X Reaction Buffer:** 0.5 M Tris-acetate (pH 7.5), 1.5 M potassium acetate, 100 mM magnesium acetate, 40 mM spermidine.

**Note:** *This buffer contains Mg<sup>2+</sup>. Do not use for the production of Transposomes.*

**EZ-Tn5 10X Stop Solution:** 1% SDS.

**Note:** *This product is not compatible with Nextera™ sequencing.*

## 4. Applications

### 1. *In Vitro* Transposon Insertion Reaction

This reaction inserts an EZ-Tn5 Transposon into target DNA, *in vitro*. The target DNA should be free of contaminating chromosomal DNA which is a direct competitor of the target DNA for insertion. Reaction conditions given have been optimized to maximize transposition frequency while minimizing multiple insertion events. Be sure to calculate the moles of target DNA used in the reaction and add an equimolar amount of the EZ-Tn5 Transposon.

1. Prepare the transposon insertion reaction mixture by adding in the following order:

1	μl	EZ-Tn5 10X Reaction Buffer
0.2	μg	target DNA**
x	μl	molar equivalent EZ-Tn5 Transposon
x	μl	sterile water to a reaction volume of 9 μl
1	μl	EZ-Tn5 Transposase
<hr/>		
10	μl	Total reaction volume

2. Incubate the reaction mixture for 2 hours at 37°C.
3. Stop the reaction by adding 1 μl EZ-Tn5 10X Stop Solution. Mix and heat for 10 minutes at 70°C.
4. Use 1 μl for electroporation into a competent bacterial strain and plate on selective media as dictated by the transposon insert. Use of a *recA*<sup>-</sup>, *endA*<sup>-</sup> strain is preferable, for target stability and subsequent purification steps (e.g. Epicentre's TransforMax™ EC100™ Electrocompetent *E. coli*\*), but not absolutely necessary. Store unused reaction mixture at -20°C.

\*\* Calculation of μmol target DNA:

$$\mu\text{mol target DNA} = \mu\text{g target DNA} / [(\# \text{ base pairs in target DNA}) \times 660]$$

For example: 0.2 μg of a 6,100 bp target clone

$$= 0.2 \mu\text{g} / [6,100 \text{ bp} \times 660] = 0.05 \times 10^{-6} \mu\text{mol} = 0.05 \text{ pmol}$$

## 2. Creating a Custom EZ-Tn5 Transposon with any DNA insert


A custom EZ-Tn5 Transposon consisting of any DNA sequence flanked by the 19-bp (ME) EZ-Tn5 Transposase recognition sequences can be generated by PCR using primers with the ME ends.

The DNA of interest, which can be anything - a resistance marker, a gene, a control element, etc. - must be amplified by PCR using primers that, in addition to having 3'-sequences homologous to the template, also have non-homologous tails with 19-base ME sequences at their 5'-ends. PCR amplification of the template using these ME-Tailed primers produces an EZ-Tn5 Transposon that can be used directly, without further purification, for in vitro insertion into any DNA target.

Design your forward and reverse primers as shown in the figures below, adding the ME sequence to the 5' end of **both** the forward and reverse primer.

ME sequence: 5'-CTG TCT CTT ATA CAC ATC T - 3'

For optimum transposition efficiency, add a 5' phosphate to both PCR primers.



## 5. Production of EZ-Tn5 Transposomes

Production of stable EZ-Tn5 Transposomes can only be accomplished in the absence of Mg<sup>2+</sup>. Do not use the EZ-Tn5 10X Reaction Buffer provided with the EZ-Tn5 Transposase to prepare EZ-Tn5 Transposomes.

- Prepare the transposome reaction mixture by adding in the following order: <sup>\*</sup>

2	μl	EZ-Tn5 Transposon DNA (100 μg/ml in TE Buffer
		[10 mM Tris-HCl (pH 7.5), 1 mM EDTA])
4	μl	EZ-Tn5 Transposase
2	μl	100% glycerol
<hr style="width: 100%;"/>		
8	μl	Total reaction volume
- Mix by vortexing. Incubate for 30 minutes at room temperature.
- Store the solution at –20°C.  
The solution will not freeze stored at –20°C and is stable for at least one year.
- Use 1 μl of the EZ-Tn5 Transposome for electroporation into a competent bacterial strain and plate on selective media as dictated by the transposon insert.

## 6. References

*Cited:*

- Goryshin, I. Y. and Reznikoff, W. S. (1998) *J. Biol. Chem.* **273**, 7367.

*Related:*

- York, D. et al., (1998) *Nucl. Acids Res.* **26**, 1927.

<sup>\*</sup> The EZ-Tn5 Transposome production protocol can be scaled up or scaled down as needed.

*These products are intended for research use only.*

*EZ-Tn5™ Transposon Tools for in vitro transposon insertion are covered by U.S. Patent Nos. 5,925,545; 5,948,622; 5,965,443, and 6,437,109; European Patent No. 0927258, and other patents issued or pending, exclusively licensed or assigned to Epicentre. These products are accompanied by a limited non-exclusive license for the purchaser to use the purchased product(s) solely for in vitro transposon insertion for life science research. Purchase of these products does not grant rights to: (1) offer products, components of products, or any derivatives thereof for resale; or (2) to distribute or transfer the products, components of products, or any derivatives thereof to third parties. Contact Epicentre for information on licenses for uses other than life science research.*

*EC100, EZ-Tn5, and TransforMax are trademarks of Lucigen Corporation.*

*Transposome is a trademark of Epicentre, Madison, Wisconsin.*

*Triton is a registered trademark of Rohm & Haas, Philadelphia, Pennsylvania.*